Every FIGHTER needs support in their corner



See how you can be there for your loved one with returning Parkinson's symptoms

Actor portrayals.

INBRIJA treats OFF periods in adults taking carbidopa/levodopa (CD/LD). INBRIJA doesn't replace CD/LD.

INBRIJA® (levodopa inhalation powder)

IMPORTANT CONSUMER SAFETY INFORMATION

What is INBRIJA?
INBRIJA is an inhaled levodopa prescription medicine used to treat the return of Parkinson's symptoms (known as OFF episodes) in people with Parkinson's disease who are treated with carbidopa/levodopa medicines. INBRIJA does not replace regular carbidopa/levodopa medicine. It is not known if INBRIJA is safe or effective in children.

Do not use INBRIJA if you: take or have taken a nonselective monoamine oxidase inhibitor such as phenelzine or tranylcypromine within the last 2 weeks.

Please see additional Important Safety Information throughout and see Patient Information Leaflet and Instructions For Use at www.INBRIJAFULLPI.com.

Inbrija®
(levodopa inhalation powder)
42 mg capsules

SYMPTOM RETURN can make even the simplest moments difficult



If your loved one is living with Parkinson's disease (PD), you may have noticed a lack of consistency in how their regular medication works over time. This can cause their PD symptoms to return (also known as OFF episodes). Symptoms may come back at the worst times, leading your loved one to avoid going out or making plans.

Many people with PD experience a return of symptoms between doses of their carbidopa/levodopa (CD/LD) medication. PD can slow the movement of food and pills through the gut, which may delay how quickly the medication is absorbed. This can cause it to take longer to start working—or stop working before the next dose.

That's one reason why PD symptoms can still return–even when your loved one takes their medication exactly as prescribed by their doctor.

What does the return of Parkinson's symptoms look like?

Knowing the signs can help you identify when your loved one's Parkinson's symptoms are returning.



- Tremor (shaking)
- Problems with balance
- Stiffness



- Anxiety*
- Mood changes*
- Difficulty thinking*

*INBRIJA has not been studied for non-movement related issues.

IMPORTANT CONSUMER SAFETY INFORMATION (Continued)

Before using INBRIJA, tell your healthcare provider about your medical conditions, including if you:

- have asthma, chronic obstructive pulmonary disease (COPD), or any chronic lung disease
- have daytime sleepiness, sleep disorders, sleepiness/drowsiness without warning, or use medicine that increases sleepiness, including antidepressants or antipsychotics
- have dizziness, nausea, sweating, or fainting when standing up
- have abnormal movement (dyskinesia)
- have mental health problems such as hallucinations or psychosis
- have uncontrollable urges like gambling, sexual urges, spending money, or binge eating

FIGHT BACK with INBRIJA



INBRIJA is an on-demand levodopa inhaler that helps address the return of PD symptoms while regular oral medication takes effect.

INBRIJA is:

- Fast-acting, with clinical improvements starting as early as 10 minutes*
- Inhaled as needed, in addition to regular CD/LD medication
- Lightweight, portable and easy to use

*In a large clinical trial, INBRIJA significantly improved motor symptoms in 30 minutes, with improvements starting as early as 10 minutes. Results may vary.

Nurse Educators† are here to help you and your fighter

Our Nurse Educators can support you and your loved one with additional training by phone or video and to answer questions about how to use the inhaler.

†Paid by Merz Therapeutics. Nurse Educators are not affiliated with INBRIJA prescribers and do not provide medical advice.

Learn more at inbrija.com or contact a Nurse Educator at 1-833-INBRIJA (1-833-462-7452)

YOU PLAY A VITAL ROLE

in advocating for your loved one's care.
Ask their doctor if INBRIJA may be right for them.

IMPORTANT CONSUMER SAFETY INFORMATION (Continued)

- have glaucoma
- are pregnant or plan to become pregnant. It is unknown if INBRIJA can harm your unborn baby.
- are breastfeeding or plan to breastfeed. Levodopa can pass into breastmilk and it is unknown if it can harm the baby.

Please see additional Important Safety Information on the back cover and see Patient Information Leaflet and Instructions For Use at www.INBRIJAFULLPI.com.

Inbrija®
(levodopa inhalation powder)
42 mg capsules

It can be helpful to

SHARE YOUR EXPERIENCE

as a care partner with your loved one's doctor



Answer the questions on the back of this tear-off page and bring it to their next appointment

Support your FIGHTER at their next appointment

A fighter never enters the ring without preparing first. Answer the questions below to get ready for your loved one's next doctor's appointment.

1 **Are you ever surprised by the return of their PD symptoms?**

- ◯ Always
- ◯ Sometimes
- ◯ Never

2 **How do you feel when their symptoms return?**

_____

_____

_____

3 **Have their symptoms ever returned during any of the following situations? (check all that apply)**

- ◯ When spending time with family
- ◯ While out in public (for example, in a restaurant or park)
- ◯ While at work
- ◯ On an important day (for example, a wedding or birthday)
- ◯ While joining in outdoor activities

4 **Have you ever avoided activities out of concern that their symptoms might return?**

- ◯ There are many things we won't do anymore
- ◯ We've stopped doing some things but still enjoy most of our favorite activities
- ◯ I'm worried we will have to stop some activities in the future for this reason
- ◯ We haven't stopped any activities

5 **How helpful would an on-demand treatment option be for managing their symptoms as they happen?**

- ◯ Very helpful
- ◯ Somewhat helpful
- ◯ Not helpful

08/2026 M-INB-US-0138

Inbrija®
(levodopa inhalation powder)
42 mg capsules

Scan to see how INBRIJA may help your loved one fight back

INBRIJA is an on-demand levodopa inhaler that helps address the return of PD symptoms. It can start working in as early as 10 minutes.* This can help your loved one get back to doing what they love most.

*In a large clinical trial, INBRIJA significantly improved motor symptoms in 30 minutes, with improvements starting as early as 10 minutes. Results may vary.

IMPORTANT CONSUMER SAFETY INFORMATION (CONTINUED)

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

Using INBRIJA and certain other medicines may affect each other and cause serious side effects. **Especially tell your healthcare provider if you take:**

- MAO-B inhibitors
- dopamine (D2) antagonists (including phenothiazines, butyrophenones, risperidone, metoclopramide)
- isoniazid
- iron salts or multivitamins that contain iron salts

How should I use INBRIJA?

INBRIJA is **for oral inhalation use only**. **Do not** swallow or open INBRIJA capsules. Only use INBRIJA capsules with the INBRIJA inhaler. **Do not** use the INBRIJA inhaler to take any other medicine.

Do not orally inhale more than 1 dose (2 capsules) for any OFF period. **Do not** take more than 5 doses (10 capsules) in a day.

What should I avoid while taking INBRIJA?

Do not drive, operate machinery, or do other activities until you know how INBRIJA affects you. Sleepiness and falling asleep suddenly can happen as late as a year after treatment is started.

What are the possible Side Effects of INBRIJA?

INBRIJA can cause serious side effects including:

- **falling asleep during normal daily activities** with or without warning. If you become drowsy, do not drive or do activities where you need to be alert for your safety or the safety of others. Chances of falling asleep during normal activities increases if you take medicine that cause drowsiness.
- **withdrawal-emergent hyperpyrexia and confusion** (fever, stiff muscles, or changes in breathing and heartbeat) if you suddenly stop using INBRIJA or carbidopa/levodopa, or suddenly lower your dose of carbidopa/levodopa.
- **low blood pressure** when standing up (that may or may not happen with dizziness, fainting, nausea, and sweating). Get up slowly after sitting/lying down.
- **hallucinations and other psychosis –** INBRIJA may cause or worsen seeing/hearing/believing things that are not real; confusion, disorientation, or disorganized thinking; trouble sleeping; dreaming a lot; being overly suspicious or feeling people want to harm you; acting aggressive; and feeling agitated/restless.
- **unusual uncontrollable urges** such as gambling, binge eating, shopping, and sexual urges has occurred in some people using medicine like INBRIJA.
- **uncontrolled, sudden body movements (dyskinesia)** may be caused or worsened by INBRIJA. INBRIJA may need to be stopped or other Parkinson's medicines may need to be changed.
- **bronchospasm –** people with asthma, COPD, or other lung diseases may wheeze or have difficulty breathing after inhaling INBRIJA. If this occurs, stop taking INBRIJA and seek immediate medical attention.
- **increased eye pressure** in patients with glaucoma. Your healthcare provider should monitor this.
- **changes in certain lab values** including liver tests.

The most common side effects of INBRIJA are cough, upper respiratory tract infection, nausea, and change in the color of saliva or spit. Another side effect of INBRIJA is sensation of choking right after use.

These are not all the possible side effects of INBRIJA

- Call your doctor for medical advice about side effects.
- You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

The risk information provided here is not comprehensive. To learn more:

- Talk to your health care provider or pharmacist.
- Visit www.inbrija.com to obtain the Full Prescribing Information, Patient Information, and Instructions for Use.
- Call 1-800-367-5109

© 2026 Merz Pharmaceuticals, LLC. Merz and the Merz logo are registered trademarks of Merz Pharma GmbH & Co. KGaA.INBRIJA and the INBRIJA logo are registered trademarks of Merz Pharmaceuticals, LLC.

08/2026 M-INB-US-0138

MERZ THERAPEUTICS

For Full Prescribing Information, Patient Information, and Instructions for Use, please visit www.INBRIJAFULLPI.com or scan QR code.